



University of
Zurich^{UZH}

Zurich Open Repository and
Archive

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

Mental and emotional representations of “weight loss”: free-word association networks in members of bariatric surgery-related social media communities

Gero, Daniel ; File, Bálint ; Hinrichs, Noreen ; Mueller, Matteo ; Ulbert, István ; Somogyvári, Zoltán ;
Bueter, Marco

Abstract: Background: Mindset and communication barriers may hinder the acceptance of bariatric surgery (BS) by the eligible patient population. Objectives: To improve the understanding of expectations, opinions, emotions, and attitudes toward weight loss among patients with obesity. Setting: Switzerland, Germany, Austria. Methods: Survey data collected from BS-related social media communities (n = 1482). Participants were asked to write 5 words that first came to their mind about “weight loss,” and to select 2 emotions, which best described their corresponding feelings. Demographic and obesity-related data were collected. Cognitive representations were constructed based on the co-occurrence network of associations, using validated data-driven methodology. Results: Respondents were Caucasian (98%), female (94%), aged 42.5 ± 10.1 years, current/highest lifetime body mass index = $36.9 \pm 9/50.7 \pm 8.7$ kg/m². The association network analysis revealed the following 2 cognitive modules: benefit-focused (health, attractiveness, happiness, agility) and procedure-focused (effort, diet, sport, surgery). Patients willing to undergo BS were more benefit-focused (odds ratio [OR] = 2.4, P = .02) and expressed more “hope” (OR = 142, P < .001). History of BS was associated with higher adherence to the procedure-focused module (OR = 2.3, P < .001), and with increased use of the emotions “gratitude” (OR = 107, P < .001), “pride” (OR = 15, P < .001), and decreased mention of “hope” (OR = .03, P < .001). Conclusion: Patients with obesity in our study tend to think about weight loss along 2 cognitive schemes, either emphasizing its expected benefits or focusing on the process of achieving it. Benefit-focused respondents were more likely to consider BS, and to express hope rather than gratitude or pride. Novel communication strategies may increase the acceptance of BS by incorporating weight loss-related cognitive and emotional content stemming from patients’ free associations.

DOI: <https://doi.org/10.1016/j.soard.2020.05.032>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-192077>

Journal Article

Updated Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Gero, Daniel; File, Bálint; Hinrichs, Noreen; Mueller, Matteo; Ulbert, István; Somogyvári, Zoltán; Bueter, Marco (2020). Mental and emotional representations of “weight loss”: free-word association networks in members of bariatric surgery-related social media communities. *Surgery for Obesity and Related Diseases*, 16(9):1312-1320.
DOI: <https://doi.org/10.1016/j.soard.2020.05.032>

Surgery for Obesity and Related Diseases

Mental and emotional representations of "weight loss": free-word association networks in members of bariatric surgery-related social media communities --Manuscript Draft--

Manuscript Number:	SOARD-19-485R2
Article Type:	Integrated Health Article
Keywords:	Bariatric Surgery; Weight loss; Free-word Association; Network; Obesity; Emotional valence; Mindset; Motivation; Body image dissatisfaction
Corresponding Author:	Marco Bueter University Hospital Zürich Zürich, SWITZERLAND
First Author:	Daniel Gero, MD
Order of Authors:	Daniel Gero, MD Bálint File, MSc. Noreen Hinrichs, MD Matteo Müller, MD István Ulbert, MD, PhD Zoltán Somogyvári, MSc., PhD Marco Bueter, MD, PhD
Manuscript Region of Origin:	SWITZERLAND
Abstract:	<p>Background: Mindset and communication barriers may hinder the acceptance of bariatric surgery (BS) by the eligible patient population. Objectives: To improve the understanding of expectations, opinions, emotions and attitudes toward weight-loss among patients with obesity. Setting: Switzerland, Germany, Austria. Methods: Survey data collected from BS-related social media communities (N=1482). Participants were asked to write 5 words that first came to their mind about "weight loss", and to select 2 emotions which best described their corresponding feelings. Demographic and obesity-related data were collected. Cognitive representations were constructed based on the co-occurrence network of associations, using validated data-driven methodology. Results: Respondents were Caucasian (98%), female (94%), aged 42.5±10.1 years, current/highest lifetime body mass index=36.9±9/50.7±8.7 kg/m². The association network analysis revealed two cognitive modules: benefit-focused (health, attractiveness, happiness, agility) and procedure-focused (effort, diet, sport, surgery). Patients willing to undergo BS were more benefit-focused (Odds ratio (OR)=2.4, P=0.02) and expressed more 'hope' (OR=142, P<0.001). History of BS was associated with higher adherence to the procedure-focused module (OR=2.3, P<0.001), and with increased use of the emotions 'gratitude' (OR=107, P<0.001), 'pride' (OR=15, P<0.001), and decreased mention of 'hope' (OR=0.03, P<0.001). Conclusion: Patients with obesity in our study tend to think about weight loss along two cognitive schemes, either emphasizing its expected benefits or focusing on the process of achieving it. Benefit-focused respondents were more likely to consider BS, and to express hope rather than gratitude or pride. Novel communication strategies may increase the acceptance of BS by incorporating weight loss-related cognitive and emotional content stemming from patients' free associations.</p>



To

Prof. Harvey Sugermann, MD
Prof. Raul Rosenthal, MD
Co-Editors-in-chief
Surgery for Obesity and Related Diseases

Zürich, 16th September 2019

Prof. Dr. med. Marco Bueter

**Leitender Arzt
Adipositaszentrum Zürich (AZZ)**

UniversitätsSpital Zürich
Klinik für Viszeral- &
Transplantationschirurgie
Rämistrasse 100
CH-8091 Zürich

Tel. 044 255 88 95
Fax. 044 255 89 42
Marco.bueter@usz.ch

Dear Professor Sugermann, dear Professor Rosenthal,

We herein submit our manuscript entitled *“Mental and emotional representations of “weight loss”: free-word association networks in members of bariatric surgery-related social media communities”* for publication in ‘Surgery for Obesity and Related Diseases’ as *Integrated Health Article*.

Our aim was to provide a patient-centered perspective to improve communicational strategies with bariatric candidates by mapping the mental and emotional representation of the term “weight loss” in patients with obesity and/or history of bariatric surgery. We used free word associations networks combined with a novel data-driven analytical method, developed and validated earlier by our co-authors. We disseminated the invitations for participation to our online survey within bariatric surgery related Facebook groups, leading to a large study cohort with complete answers (n = 1482). The association network analysis revealed two distinct cognitive modules: benefit-focused (most frequent associations: health, agility, attractiveness) and procedure-focused (most frequent associations: sport, diet, effort).

We believe that our findings may serve as basis to improve patient-centered communication strategies and therefore may be highly relevant for the readers of ‘Surgery for Obesity and Related Diseases’.

The study was presented in part at the 24th World Congress of the *International Federation for the Surgery of Obesity and Metabolic Disorders* in Madrid, Spain, on 5th September 2019. This manuscript has not been previously published or submitted elsewhere for publication.



The manuscript has been approved by all authors. The study was entirely supported by the University of Zürich, without any external funding. The study could not have been achieved without a multi-disciplinary multi-national collaboration between surgeons and bionic engineers, which justifies the involvement of 7 co-authors.

We would be most grateful if our manuscript was considered for publication in 'Surgery for Obesity and Related Diseases'. Please do not hesitate to contact us for further information.

Yours sincerely,

Prof. Marco Bueter, MD, PhD

SURGERY FOR OBESITY AND RELATED DISEASES (SOARD)
Author Disclosure of Relevant Financial Relationships

NAME: Marco Bueter, MD, PhD (corresponding author)

TITLE OF ARTICLE: Mental and Emotional Representations of "Weight Loss": Findings from a Survey Using Free-word Association Networks in Bariatric Surgery-related Social Media Communities.

The American Society for Metabolic and Bariatric Surgery must ensure that anyone who is in a position to control the content of the education activity has disclosed to us all relevant financial relationships with any commercial interest (see below for definitions). Please indicate below any commercial interests that would pertain to the content of the article being submitted to SOARD.

- ☒ None of the co-authors have anything to disclose (if you are the corresponding author).
- ☐ I do not have anything to disclose (if you are an individual author).
- ☒ I am disclosing all relevant financial relationships with any commercial interests that pertain to this article (see below).

- List the names of any proprietary entities producing marketing, re-selling, or distributing health care goods or services consumed by, or used on patients with which you or your spouse/partner have, or have had, a **relevant** financial relationship within the past 12 months.
- Explain what you or your spouse/partner received (ex: salary, honorarium etc).
- Specify your role.

Commercial Interest	Nature of Relevant Financial Relationship (Include all those that apply)	
	What I or spouse/partner received	My role
Johnson & Johnson	Honorarium	Speaker
Medtronic	Honorarium	Speaker

What was received: Salary, royalty, intellectual property rights, consulting fee, honoraria, ownership interest, (e.g., stocks, stock options or other ownership interest, excluding diversified mutual funds), or other financial benefit.

My Role(s): Employment, management position, independent contractor (including contracted research), consulting, speaking and teaching, membership on advisory committees or review panels, board membership, and other activities.

Through our review process, should it be determined that a bias is apparent within the content of the article, you will be requested to revise your manuscript prior to publication.

If your article describes the use of a device, product, or drug that is not FDA approved or the off-label use of an approved device, product, or drug or unapproved usage, it is your responsibility to disclose this information within the manuscript.

Glossary of Terms

Commercial Interest
A commercial interest is any entity producing, marketing, re-selling, or distributing health care goods or services consumed by, or used on, patients. The ACCME does not consider providers of clinical service directly to patients to be commercial interests.

Financial relationships
Financial relationships are those relationships in which the individual benefits by receiving a salary, royalty, intellectual property rights, consulting fee, honoraria, ownership interest (e.g., stocks, stock options or other ownership interest, excluding diversified mutual funds), or other financial benefit. Financial benefits are usually associated with roles such as employment, management position, independent contractor (including contracted research), consulting, speaking and teaching, membership on advisory committees or review panels, board membership, and other activities from which remuneration is received, or expected. ACCME considers relationships of the person involved in the CME activity to include financial relationships of a spouse or partner.

Relevant financial relationships
ACCME focuses on financial relationships with commercial interests in the 12-month period preceding the time that the individual is being asked to assume a role controlling content of the CME activity. ACCME has not set a minimal dollar amount for relationships to be significant. Inherent in any amount is the incentive to maintain or increase the value of the relationship. The ACCME defines "relevant" financial relationships" as financial relationships in any amount occurring within the past 12 months that create a conflict of interest.

Conflict of Interest
Circumstances create a conflict of interest when an individual has an opportunity to affect CME content about products or services of a commercial interest with which he/she has a financial relationship.

Checklist

Authors must submit their manuscripts to the SURGERY FOR OBESITY AND RELATED DISEASES (SOARD) Editorial Office via the Elsevier Editorial System (EES) website for this journal; go to <http://ees.elsevier.com/SOARD> and select "Submit Manuscript". All correspondence regarding submitted manuscripts will be handled via e-mail through EES. Please do not send duplicate copies of the manuscript to the editorial office.

A COMPLETED CHECKLIST FORM MUST ACCOMPANY EACH MANUSCRIPT SUBMITTED TO SOARD. **CHECK ALL ITEMS APPLICABLE TO YOUR SUBMISSION. IT IS AN ETHICAL VIOLATION TO CHECK AN ITEM WITHOUT HAVING RESPONDED TO IT.**

Full Guidelines for Authors are provided in the journal and online at: [Guide for Authors](#).

GENERAL

☒ Potential conflicts of interest, whether of a financial or other nature related to the topic of the manuscript, are disclosed by checking the appropriate box on the author agreement form and listing the potential conflicts in an attachment to the author agreement form.

In addition, the manuscript itself must include a Disclosure Statement in which every author must list any financial or personal relationships that could inappropriately bias the work.

☒ Pages are numbered consecutively

☒ I certify that the manuscript did not involve the use of animal or human subjects
OR

I certify that a statement confirming that appropriate institutional and/or ethics committee approval has been obtained and is stated in the Methods section of the manuscript.

☒ All organizations that have funded this research have been acknowledged, including grant numbers where appropriate

☒ Lines are numbered consecutively at 5 line intervals throughout the manuscript using the line numbering feature available in word processing software.

☒ The article is typed in 12-point type, double-spaced with one-inch margins

☒ Title page is uploaded separately (see below for title page requirements)

☒ No author or institutional information in the manuscript text

☒ **Article is written in American English (e.g. anesthesia rather than anaesthesia)**

☒ Unless a multicenter study, there are no more than 6 co-authors on an Original Manuscript and no more than 3 on a Case Report. Your article will be automatically rejected without review unless you have **VERY STRONG EVIDENCE** that more than 6 authors are necessary.

☒ Only the following abbreviations will be accepted: RYGB or LRYGB for Roux-en-Y Gastric Bypass or Laparoscopic RYGB; BPD for Biliopancreatic Diversion and for BPD with Duodenal Switch, BPD/DS; LAGB for laparoscopic adjustable gastric band; SG for Sleeve Gastrectomy

☒ ALL abbreviations (including BMI) are defined when first presented in both the abstract and the manuscript

☒ Weight loss must be expressed as change in BMI, **AS WELL AS** % Excess Weight Loss (%EWL), with the calculation of ideal body weight as that equivalent to a BMI of 25 kg/m² and/or % Excess BMI Lost (%EBMIL) with excess BMI > 25 kg/m² **AS WELL AS** % total body weight loss

☒ Data extending beyond 30 days **MUST INCLUDE** lost to follow-up information in the Abstract and Results section, including all tables and figures, with the denominator provided as to how many patients were available at **EACH TIME POINT** and the number of patients actually seen

☒ Each manuscript requires a Conclusions paragraph. Prior to this paragraph, there needs to be a

presentation of the limitations of the study

☒ Following the Conclusion paragraph there needs to be a disclosure of all conflicts of interest or, if none, so state.

☒ Studies in humans require informed consent and approval from the local Institutional Review Board (IRB) and in animals from the Animal Care and Use Committee and is addressed in the Materials and Methods section.

☒ Permission to reprint material from another source in print and electronic form has been obtained.

☒ If a clinical trial, it was registered in the National Institutes of Health web site: www.clinicaltrials.gov or one of the equivalent web sites such as: <http://www.anzctr.org.au>; <http://www.clinicaltrials.gov>; <http://isrctn.org>; <http://www.trialregister.nl/trialreg/index.asp>; or <http://www.unin.ac.jp/ctr>. This information has been stated in both the Cover Letter and the Materials and Methods section of the manuscript.

☒ For randomized controlled trials, a completed CONSORT checklist accompanies the manuscript. Download at: http://www.consort-statement.org/mod_product/uploads/CONSORT%202001%20checklist.pdf

☒ For studies of diagnostic accuracy, a completed STARD checklist accompanies the manuscript. Download at: http://www.stard-statement.org/checklist_maintext.htm

☒ For meta-analyses and systematic reviews of randomized controlled trials, the PRISMA guidelines are followed. Download at: <http://www.prisma-statement.org/2.1.2%20-%20PRISMA%202009%20Checklist.pdf>

☒ For meta-analyses and systematic reviews of observational studies, the MOOSE guidelines are followed. Download at: <http://www.editorialmanager.com/jognn/account/MOOSE.pdf>

☒ For observational studies, a completed STROBE checklist accompanies the manuscript. Download at:

COVER LETTER

☒ A cover letter is submitted indicating the intent to submit to *Surgery for Obesity and Related Diseases*.

☒ Publication elsewhere (including electronic media) of any of the material in the manuscript submitted, other than an abstract of not more than 300 words, is identified and a copy of the other publication is submitted to the editorial office.

☒ If a version of the manuscript has previously been submitted for publication to *Surgery for Obesity and Related Disease*, comments from the peer reviewers and an indication of how the authors have responded to these comments are included.

TITLE PAGE

☒ The title, author name(s) and major degree(s), affiliation(s), and the source(s) of funding for the work or study are provided.

☒ The name, address, telephone and fax numbers, and e-mail address of the corresponding author are given.

☒ The manuscript title is no longer than 150 characters (letters and spaces) and does not contain any abbreviations or acronyms.

☒ A short title (no more than 50 characters) is provided at the bottom of the page for use as a running foot.

☒ The only acknowledgments are of financial or other substantive assistance. All individuals named in the acknowledgments have given written permission to be named.

ABSTRACT (for original and review articles only)

☒ For original research reports and review articles, the abstract is limited to 250 and 300 words, respectively, and in the case of original articles follows the appropriate structured abstract format. **Review articles do not require a structured format.** Abstracts are not used in Case reports, Controversial Issues, Surgeon at Work and Updates.

☒ If the research identified important variables that lacked statistical difference, the study's power to detect

such difference is addressed in the abstract.

☒ Footnotes, references, and commercial names are not used in the abstract.

☒ The abstract precedes the manuscript document.

☒ Keywords are added at the end.

REFERENCES

☒ References correspond to the specifications of the "Uniform Requirements for Manuscripts Submitted to Biomedical Journals." Examples of specific types of references are available at:

http://www.elsevier.com/wps/find/journaldescription.cws_home/703772/authorinstructions

☒ References are identified on the line within parentheses **in superscript**, e.g., Schauer et al. ⁽³⁾.

☒ Each reference is cited in the text.

☒ References are double-spaced and are **numbered consecutively** in the order in which they appear in the text.

☒ ALL authors have been listed when 6 or less. Only when more than 6, should the first 3 authors be listed followed by "et al."

☒ Unpublished data, personal communications, submitted manuscripts, statistical programs, papers presented at meetings, and non-peer-review publications are not listed in the reference list.

TABLES

☒ Each table is on a separate sheet of paper with its title and uploaded separately from the manuscript.

☒ Tables are numbered with Arabic numerals, e.g. Table 1, Table 2.

☒ Each table contains all necessary information in order that it may stand alone, independent of the text, including definitions of all abbreviations in the table in a legend beneath the table.

☒ No table repeats all data presented in the text.

FIGURES

☒ Digital art guidelines and artwork checklist on the journal's web site (http://www.elsevier.com/framework_authors/Artwork/Artwork_2010.pdf) have been consulted.

☒ Each figure is cited in the text.

☒ Figures have a minimum width of 3¼ inches and height of 2 inches.

☒ Images are submitted in TIFF (Tagged Image File Format) or EPS (Encapsulated PostScript) formats

☒ Letters and identifying marks (e.g. arrows) are clear and sharp, and the critical areas of radiographs and photomicrographs are identified. **All figures and edited tables use at least 300 DPI**

☒ **Figures and tables are of high quality and sufficient size and clarity for reproduction in the journal**

☒ All text that may identify a patient, including initials of the patient's name) is removed.

☒ Explanatory material appears in the accompanying legend and not on the figure itself, including definitions of all abbreviations in the figure in a legend beneath the table.

☒ Legends are listed on one page at the end of the manuscript and must have definitions for all abbreviations.

☒ Each figure is uploaded separately from the manuscript

☒ I understand that color figures are welcome and will appear in color on the web at no extra charge; however, there is a charge for the reproduction of color illustrations in print (\$650 for the first color figure and \$100 for each additional color figure) unless you request an exception and it is waived by the Editor and Publisher. Alternatively, the illustrations can be reproduced in print in black and white at no extra charge. Be sure that charts and graphs have been designed with this in mind so that the data are clear when published in black and white, i.e. different formats for the different columns (clear, hash marks, stippled, etc.) or graphic lines (solid, dashed, dotted, etc.).

☐ I plan on paying for color images

Reviewer Suggestions

Manuscript

Gero, File et al. Mental and emotional representations of “weight loss”: free-word association networks in members of bariatric surgery-related social media communities. *Surgery for Obesity and Related Diseases*. 2019

Reviewer Suggestion #1

Prof. Dr. **Sivamainthan Vithiananthan** FACS FASMBS FICS

Chief of minimally invasive and bariatric surgery at the Center for Bariatric Surgery, Alpert Medical School - Brown University, Providence, RI

Address: 195 Collyer St suite 302, Providence, RI 02904, USA

Phone: +1 401-793-5140

Email: svithiananthan@Lifespan.org

Reviewer Suggestion #2

Prof. Dr. Mirto Foletto

Chief of Bariatric Surgery

Address: CLINICA CHIRURGICA 1, University of Padova, Via Giustiniani, 2, 35128 PADOVA, Italy

Phone: +39 840-000664

Email: mirto.foletto@gmail.com

Reviewer Suggestion #3

Prof Antonio Torres, M.D., Ph.D., F.A.C.S., FASMBS

Chair Of Surgery at Complutense University of Madrid

Pza. Ramón y Cajal, s/n

Ciudad Universitaria

28040 - MADRID

Email: ajtorresgarcia@gmail.com

Reviewer Suggestion #4

Univ.Do. Dr. Gerhard Prager

Chair of Bariatric Surgery

Medizinische Universität Wien

Universitätsklinik für Chirurgie

Währinger Gürtel 18-20

1090 Wien

Email: Gerhard.prager@meduniwien.ac.at

Point to Point reply

Ms No.: SOARD-19-485R1

Title: Mental and emotional representations of "weight loss": free-word association networks in members of bariatric surgery-related social media communities

We would like to thank the editor and both reviewers for their interest and critical evaluation of our manuscript. Please find below our point-to-point replies.

We remain available for any further questions or comments.

Associate Editor

Comment #1: The respondents are 98% Caucasian and 94% female yet the manuscript's conclusions focus on attitudes of people with obesity. The findings really can only speak to the thoughts/attitudes of White women with obesity and this should be strongly tempered throughout.

Our reply: This is now emphasized in the first paragraph of the Discussion, as well as in the Limitations. We also updated the Supplemental Table 1, which now contains detailed demographic comparison of all subgroups.

Comment #2: Please add more about the demographic questions. How was BMI determined?

Our reply: We added the following paragraph to Methods:

"Basic Demographics

Respondents were asked to self-report their age, sex, ethnicity, civil status, height, maximal and current body weight, past or planned bariatric operations, previous non-surgical weight loss attempts and their social interactions with bariatric patients. Body mass index (BMI) was calculated by the investigators based on the anthropometric data provided by the participants."

Comment #3: How were the basic emotions selected?

Our reply: The selection of the emotions was largely built on the ten basic emotions of Izard (Izard, 1977) and the eleven pairs of positive and negative emotion pairs of Robinson (Robinson, 2008 – this reference has now been added). Respondents could choose two out of the 20 emotions to label for each of their own free word associations. Although each emotional label had an opposite pair, the labels did not appear as opposites in the questionnaire. The same emotional labeling process has been applied successfully in our recent validation study (File et al, 2019, doi: 10.3758/s13428-018-1090-z).

Comment #4: Reviewer 2 questioned the validity of the Figure Rating Scales in this population and more information should be provided. These figures likely have a ceiling effect in this population and a full range of scores aren't possible. Please spell out BID when first discussed. A discrepancy between a current and ideal body size may not be indicative of BID in this population as it may reflect current BMI versus health-related expectations. Indeed using a discrepancy score using these figures for BID came from the literature in normal weight women and those with more traditional eating disorders (e.g., anorexia). Similarly, the finding that the discrepancy is greater in those with higher BMI's seems rather tautological.

Our reply: Thank you for this important comment, which we now mention in the Methods. Although developed for the "normal weight" general population, the Stunkard rating Scale has been regularly used in the context of obesity / bariatric surgery. A study published earlier in SOARD has also used the Stunkard scale:

Ratcliff MB, Eshleman KE, Reiter-Purtill J, Zeller MH. Prospective changes in body image dissatisfaction among adolescent bariatric patients: the importance of body size estimation. Surg Obes Relat Dis. 2012;8:470-5 (this reference has now been added).

Indeed, BID score in the bariatric population may reflect current BMI versus health-related expectations. We found a moderate correlation ($r = 0.57$) between current BMI and the BID score. To better investigate the Editor's concern, we now correlated health related expectation (expected body image) and current body image with the current BMI value and updated the Results accordingly:

"Overall, current BMI showed a) a strong correlation with BID (Pearson's $r = 0.6$, $P < 0.001$, Supplemental Fig. 4), b) an even stronger correlation with current body image ($r = 0.77$, $P < 0.001$) and c) a weaker correlation with the expected body image ($r = 0.42$, $P < 0.001$)."

According to these results, health related expectations show a lower correlation with the BMI than the current body image. We conclude that current and expected body image have a significant relationship with the current BMI, but to a different extent.

Comment #5: I'm curious about the subset who is "not interested in BS". They seem unusual since if they were truly uninterested why would they be in social medial groups for bariatric surgery? More discussion of these individuals is warranted.

Our reply: We added the following subsection on this subgroup in Results:

"Respondents who would not consider to undergo BS

Out of the 409 respondents without prior BS, 66 stated to have no interest to undergo BS. This subgroup had a mean age of 43.6 years and consisted of Caucasian (91%) females (92%) with a BID of 2.2 who managed to lose weight non-surgically (highest lifetime BMI: 45.4±8.3 kg/m², current BMI: 37.9±8.8 kg/m²). Only 53% of this group had regular contact with BS patients."

and to Discussion:

"A subgroup of the respondents stated not to be interested to undergo BS, nevertheless, by following BS-related social media they manifested interest towards BS and seem to be seeking facts and/or patient narratives to assist their health-related decisions. "

Comment #6: The completion rate is noted as 67.1% but those are individuals who went to the landing page. How many people visit these social media sites?

Our reply: This question is hard to answer because we do not have exact data. As stated in Methods, the Adipositas Zürich website has 4100 followers and the Adipositas Chirurgie - Fragen und Antworten' has 1500. We encouraged visitors to share the link of the questionnaires within their own networks. The original invitation to participate on the Facebook wall of the Adipositas Zurich group was shared 113 times. We believe that it is accurate to report the completion rate only among those who actively clicked on the invitation and started to read the questionnaire on the landing page.

Comment #7: Please use a term like sub-optimal outcome rather than "failure" when discussing weight loss as it is less pejorative

Our reply: : "Failure" was used in relation with weight loss in two sentences. These two sentences have been rephrased as follows:

"According to a Brazilian study in patients experiencing postbariatric weight regain, the feeling of gratitude toward BS remained, whereas pride was replaced by a feeling of sub-optimal outcome and low self-esteem [29]."

"However, unrealistic expectations can be responsible for the cycle of sub-optimal outcomes and renewed efforts to achieve self-change (i.e.: false-hope syndrome, "yo-yo effect")[24]."

REVIEWER 1

Comment #1: Line 52 - suggest "incorrect" instead of "wrong" information

Our reply: Corrected.

Comment #2: Line 77 - suggest inclusion of adults with obesity "seeking or post surgery" since there is a whole group not interested/seeking surgery that are not included here

Our reply: The sentence has been corrected: *"We conducted an anonymized online survey to understand how adults with severe obesity (those who seek BS, postbariatric patients, and those who are not interested to undergo BS) perceive weight loss."*

Comment #3: Line 100 - successfully applied 'earlier' (I think you mean in earlier studies, or in prior work?)

Our reply: In "a recent study" - we have now rephrased this sentence.

REVIEWER 2

Comment #1: Line 40: the first line seems strongly worded - "Patients with obesity tend to think..." Can this be tempered a bit? (e.g., Patients with obesity in our study...)

Our reply: Done.

Comment #2: Line 54: is this suboptimal cooperation or suboptimal communication between primary care physicians and bariatric centers? Just clarifying!

Our reply: This sentence has been rephrased to:

"Mindset and communication barriers seem to influence the diffusion of BS: lack of information, incorrect information, and stigmatization of individuals with obesity have proven to contribute to sub-optimal use of health care by the eligible patients and to insufficient cooperation between primary care physicians and BS centers [3, 4]."

Comment #3: Body Image Dissatisfaction: Has the Stunkard Figure Rating Scale been validated in the use of people with obesity? I am also unclear about how the analysis for BID related to the overall aims of the study.

Our reply: Please read our reply to Associate Editor's Comment#4 on the use of the Stunkard scale in the context of obesity. We hypothesized that BID may influence the cognitive representation of weight loss, however, as shown in Table 1, the BID scores were similar between the Benefit-focused and Procedure-focused groups. Body image is an important indicator of patients' wellbeing after BS [ref#27] and was higher in participants who planned to undergo BS, compared to those who did not, or already had BS (excerpt from Discussion).

Comment #4: Are there any a priori hypotheses?

Our reply: This is the first time we used social media to recruit respondents for a bariatric surgery related study. We had troubles to estimate the expected cohort size and demographic composition of respondents, making it difficult to draft a priori hypotheses. Additionally, the free word association technique is a data-driven explorative analysis. It has the advantage to explore cognitive representations beyond preconceptions, but it is almost impossible to guess the number and content of the evolving semantic groups. Therefore, our hypotheses were only related to the emotional labels of the associations.

The following amendments have been added to the Introduction, Results and Discussion:

“We hypothesized that BS had a “positive” impact on the emotional labeling of *weight loss* related free word associations.”

“Postbariatric patients had a significantly higher proportion of positive emotional labels than non-operated adults with obesity (79% vs 75%, $t(1480)=1.82$; $P = 0.002$, $d=0.18$). “

“In line with our hypothesis, history of BS favored the use of positive emotional labels, however, the effect size was rather small. In fact, all respondents selected mainly positive emotions to label their associations. Remarkably, patients willing to undergo BS were more likely to express *hope*, while patients after BS expressed *gratitude* and *pride* more frequently, suggesting that BS may fundamentally change the individuals’ affective connection with weight loss.”

Comment #5: Line 100/101: Are there any other published studies to use this method for emotional labeling?

Our reply: Yes, our group recently published a methodological validation study: File et al, 2019, doi: 10.3758/s13428-018-1090-z. Please see our reply to the Associate Editor’s Comment#3.

Comment #6: Consider demographic tables for the following:

- a. $\geq / < 35$ BMI
- b. With interest for BS vs no interest
- c. With BS vs without

These may provide clarification for the existing Table 2a and 2b

Our reply: Thank you for this comment. This information is available in the Supplemental Table 1, which has been updated in this new version to fully respond to the Reviewer’s comment.

Comment #7: Table 2A: were there any demographic differences between the two groups (Respondents with \geq Class Obesity and Respondents with prior bariatric surgery)?

Our reply: Demographic differences between subgroups are shown in Supplemental Table 1. “Respondents with \geq Class II Obesity” and “Respondents with prior bariatric surgery” are subgroups of the entire cohort containing 408 overlapping individuals (those who had \geq Class II Obesity and also had prior bariatric surgery). In the first logistic regression of the “old” Table 2A, the reference was the subgroup of respondents <Class II Obesity, while in the second one, those without prior BS. We re-organized this Table and its legends to increase clarity.

Comment #8: Are there any more clinical implications beyond (Lines 255-258)?

Our reply: We updated this section, which now reads as:

“...this study provided a comprehensive structure of adults’ with obesity free expressions about weight loss, enhancing our understanding of patients’ priorities and contributing to the improvement of health care-related communicational schemes. Discrepant expectations may significantly increase postoperative dissatisfaction with body image and decrease the extent of achievable weight loss [33], therefore setting realistic preoperative expectations and stepwise postoperative goals are crucial. Our findings also promote the collaborative approach between healthcare providers to support evidence-based treatments of obesity [34].”

Other applied implications are mentioned later in the same paragraph:

“Consequently, the network-based map of associations to weight loss presented in this study may serve as a navigational tool for caregivers, policy actors and researchers to better identify patients’ implicit interest toward weight loss and to design new interventions aiming to tackle mindset barriers in the acceptance of BS among patients with obesity [38].”

“...the relevant positioning of “weight loss” can influence large groups of people with obesity and change social norms [39]. Once a mindset has been primed or activated, there is an increased likelihood that it will be used in upcoming tasks to interpret new information and to define subsequent actions [24].”

Comment #9: If this group is looking to expand on this work, consider examining cognitive schemas in primary care providers; I wonder how this may affect referrals!

Our reply: Thank you, we fully agree and already started to brainstorm on the next steps.

Title

Mental and emotional representations of “weight loss”: free-word association networks in members of bariatric surgery-related social media communities

Authors

Daniel Gero, MD^{1*}, Bálint File, MSc^{2,3*}, Noreen Hinrichs, MD¹, Matteo Mueller, MD¹, István Ulbert, MD, PhD², Zoltán Somogyvári, MSc PhD³, Marco Bueter, MD, PhD¹

*DG and BF contributed equally

¹ Department of Surgery and Transplantation, University Hospital Zurich, Switzerland

² Faculty of Information Technology and Bionics, Pázmány Péter Catholic University, Budapest, Hungary

³ Wigner Research Centre for Physics, Hungarian Academy of Sciences, Budapest, Hungary

Manuscript category: Integrated Health Article

Communication: The study was presented in part at the 24th International Federation for the Surgery of Obesity and Metabolic Disorders World Congress on 5th September 2019 in Madrid, Spain, and at the 106th Annual meeting of the Swiss Society of Surgeons on 17 May 2019 in Bern, Switzerland.

Funding: This work was supported by the assistant-professorship research grant awarded by the University of Zurich to Prof. Dr. med. Marco Bueter, PhD; Bálint File, MSc was supported by the ÚNKP-18-3-III-PPKE-65 New National Excellence Program of the Hungarian Ministry of Human Capacities.

Correspondence to:

Prof. Dr. med. Marco Bueter, PhD

Department of Surgery and Transplantation, University Hospital Zurich, Switzerland

Rämistrasse 100, 8006 Zurich, Switzerland

Phone: +41 (0)44 255 8895

Fax: +41 (0)44 255 8941

Email: marco.bueter@usz.ch

Running title: Mental and emotional representations of weight loss

Title

Mental and emotional representations of “weight loss”: free-word association networks in members of bariatric surgery-related social media communities

5 Authors

Blinded

Manuscript category: Integrated Health Article

10 **Communication:** The study was presented in part at the 24th International Federation for the Surgery of Obesity and Metabolic Disorders World Congress on 5 September 2019 in Madrid, Spain and at the 106th Annual meeting of the Swiss Society of Surgeons on 17 May 2019 in Bern, Switzerland.

Funding: This work was supported by the assistant-professorship research grant awarded by the **Blinded** to Prof. Dr. med. **Blinded**, PhD; **Blinded**, MSc was supported by the **Blinded** New
15 National Excellence Program of the **Blinded**.

Correspondence to:

Blinded

Running title: Mental and emotional representations of weight loss

20 **Abstract**

Background: Mindset and communication barriers may hinder the acceptance of bariatric surgery (BS) by the eligible patient population.

Objectives: To improve the understanding of expectations, opinions, emotions and attitudes toward weight-loss among patients with obesity.

25 **Setting:** Switzerland, Germany, Austria.

Methods: Survey data collected from BS-related social media communities (N=1482). Participants were asked to write 5 words that first came to their mind about “weight loss”, and to select 2 emotions which best described their corresponding feelings. Demographic and obesity-related data were collected. Cognitive representations were constructed based on the co-occurrence network of
30 associations, using validated data-driven methodology.

Results: Respondents were Caucasian (98%), female (94%), aged 42.5 ± 10.1 years, current/highest lifetime body mass index = $36.9 \pm 9 / 50.7 \pm 8.7$ kg/m². The association network analysis revealed two cognitive modules: benefit-focused (health, attractiveness, happiness, agility) and procedure-focused (effort, diet, sport, surgery). Patients willing to undergo BS were more benefit-focused (Odds ratio (OR)=2.4, P=0.02) and expressed more ‘hope’ (OR=142, P<0.001). History of BS was associated with
35 higher adherence to the procedure-focused module (OR=2.3, P<0.001), and with increased use of the emotions ‘gratitude’ (OR=107, P<0.001), ‘pride’ (OR=15, P<0.001), and decreased mention of ‘hope’ (OR=0.03, P<0.001).

Conclusion: Patients with obesity in our study tend to think about weight loss along two cognitive
40 schemes, either emphasizing its expected benefits or focusing on the process of achieving it. Benefit-focused respondents were more likely to consider BS, and to express hope rather than gratitude or pride. Novel communication strategies may increase the acceptance of BS by incorporating weight loss-related cognitive and emotional content stemming from patients’ free associations.

Keywords

45 Bariatric Surgery; Weight loss; Free-word Association; Network; Obesity; Emotional valence; Mindset; Motivation; Body image dissatisfaction

Introduction

Bariatric surgery (BS) is currently the most effective modality in the treatment of severe obesity and related diseases [1]. Despite its growing worldwide popularity as a treatment option, BS is rather underused by the eligible patient population [2]. Mindset and communication barriers seem to influence the diffusion of BS: lack of information, ~~wrong~~ incorrect information, and stigmatization of individuals with obesity have proven to contribute to sub-optimal use of health care by the eligible patients and to insufficient cooperation between primary care physicians and BS centers [3, 4]. ~~Suboptimal cooperation between primary care physicians and bariatric centers~~ This may also hinder the identification and referral of eligible patients and undermine optimal postbariatric follow-up [4].

Available literature on motivational drivers in favor of seeking BS remains scarce and stems mainly from small cohorts [5-7]. Nevertheless, corroborative evidence shows that the willingness to improve health is the cornerstone for seeking BS, contextualized with psychological, emotional, environmental and social factors [8, 9]. Within the developing framework of person-centered care, the patients' narrative is a key factor in improving decision-making [10]. However, due to the sensitive, personal and private nature of patients' perception of their own body weight, direct questions from healthcare professional may be perceived as judgmental or scolding, and may rapidly close down communication [11]. In this sensitive context, an anonymized online survey with a free word association task seems to be an optimal approach to scope group opinion [12].

Free word association is a widely used technique in market research and psychology to encourage respondents to express openly their underlying motivations, beliefs, attitudes or feelings regarding a specific topic [13]. This technique enhances the unconstrained expressions of respondents and overcomes limitations of predefined questionnaires.

We aimed to map the mental and emotional representation of the term *weight loss* in patients with obesity who had a documented interest in BS, using free word associations combined with a novel data-driven associative analytical method [12]. Further, we aimed to investigate the relationship between these representations and clinical parameters, such as stage of obesity and previous or planned experience with BS. We hypothesized that BS had a “positive” impact on the emotional labeling of *weight loss* related free word associations. Our findings may serve as basis to improve patient-centered communication strategies, online and in clinical practice alike [14].

Methods

We conducted an anonymized online survey to understand how adults with severe obesity (those who seek BS, postbariatric patients, and those who are not interested to undergo BS) perceive *weight loss*.~~We conducted an anonymized online survey to understand how adults with obesity perceive *weight loss*.~~ To ensure that participants are familiar with BS, respondents were recruited on a voluntary basis from bariatric surgery-related German-speaking social media groups between 02/11/2018 and 04/04/2018. An invitation for participation and three reminders every two weeks were posted on the ‘Adipositas Zürich’ (4100 followers) and ‘Adipositas Chirurgie - Fragen und Antworten’ (1500 followers) Facebook© groups, and respondents were encouraged to share the invitation within their own social networks. Thus, the survey was “open” to any respondents with a presumed interest in BS. We aimed to enroll a convenience sample of >500 respondents. The survey was administered via Typeform™ (Barcelona, Spain) (Supplemental Methods 1.), and only complete questionnaires could be submitted. Respondents could not be backtracked to a specific Facebook© profile. The Cantonal Ethics Committee of Zurich concluded that this study did not fall under the scope

of the Swiss Human Research Act, thus ethical approval was not required. The manuscript was prepared according to the Checklist for Reporting Results of Internet E-Surveys (CHERRIES) guidelines [15]. Network analyses were performed using Matlab 2017 (The Mathworks); logistic regressions and correlations were performed using R Software for Statistical Computing v3.5.2 (R Foundation, Vienna, Austria).

Building the dataset

Basic Demographics

Respondents were asked to self-report their age, sex, ethnicity, civil status, height, maximal and current body weight, past or planned bariatric operations, previous non-surgical weight loss attempts and their social interactions with bariatric patients. Body mass index (BMI) was calculated by the investigators based on the anthropometric data provided by the participants.

Free word associations

~~In addition to demographic questions, p~~Participants were asked to write five words, which first came to their mind about “weight loss”. Further, they had to select two emotions from a list of 20 basic emotions (interest, anxiety, empathy, contempt, surprise, indifference, hope, fear, gratitude, anger, joy, sadness, calmness, frustration, pride, shame, generosity, envy, sympathy, antipathy) that best described their feelings about each of the five associations they provided [16, 17]. Emotional labeling was successfully applied ~~earlier in a recent study~~ to improve the interpretation of free-word associations by the addition of affective content [12].

Body image dissatisfaction (BID)

The color version of the reliable and validated Stunkard Figure Rating Scale was used to assess current and ideal body size estimations [18]. This instrument uses a series of 9 gender-specific silhouettes that gradually increase in body size. Although developed for normal weight adults with traditional eating disorders (e.g.: anorexia nervosa), this scale has been previously applied in the context of BS [19]. Respondents were instructed to select the figure that indicates (a) how they look currently and (b) how they would like to look like. A discrepancy score indicating Body Image Dissatisfaction (BID) was calculated by subtracting scores of the ideal figure (b) from the current (a) figure. Lower discrepancy scores were interpreted as lower body image dissatisfaction.

Data pre-processing

Response rate was calculated based on definitions and metrics proposed by the American Association for Public Opinion Research [20]. Respondents who provided incoherent data (i.e.: highest lifetime ~~body mass index~~ (BMI) < current BMI) or were found to be outside the target group (i.e.: highest lifetime BMI < 35 kg/m², history of gastric banding, etc.) were excluded from the study cohort in a step-wise manner (Supplemental Fig. 1). Free word associations to “weight loss” were first spellchecked and lemmatized (i.e.: to return the base of a word) by two independent native German speakers. Associations were merged if their English translation was identical. Idiosyncratic expressions (associations with <9 occurrences) were left out.

Association network analysis

In order to extract the most prominent mental representations from the numerous individual associations provided by the respondents, we applied an in-house developed algorithm, using a network-based methodology validated and described earlier by our group [12]. To create such a

network, nodes (free word associations) and edges (statistical co-occurrences between two associations) were determined. Statistical co-occurrence was calculated by log-likelihood ratios (LLR). The LLR value can denote attractive (when two associations are often mentioned together) and repulsive connections (when two associations are rarely mentioned together) with a corresponding level of significance. In the network of statistically related associations, modularity maximization based consensus clustering algorithm was applied to explore any densely connected subnetworks (i.e.: modules) [21]. We have previously demonstrated that these modules of associations - as consistent patterns in individual association sequences – are able to capture the most prominent cognitive representation of a given cue [12]. Associations with weak module attachment were separately identified, since they may serve as bridges in future communicational strategies to simultaneously address individuals with different cognitive modular membership.

Qualitative and quantitative analysis of the map

First, we used descriptive statistics to characterize the subgroups of interests, as well as the members of the two cognitive modules. *Second*, we created explorative models by logistic regression to describe the relationship between free-word associations (cognitive module membership, emotional labels) and clinical parameters (stage of obesity, BS status, interest to undergo BS). The cognitive module of respondents was determined by the association cluster to which the majority of their associations belonged. Emotions were expressed as frequency (%) and the unfrequently (<4%) mentioned emotions were not included in the models to decrease false discovery rate. Respondents' current BMI status was dichotomized according to World Health Organisation obesity class II (\geq / $<$ 35 kg/m²), since this is the currently recommended cutoff for being eligible to BS [1]. We also created subgroups based on the BS status of respondents: “underwent BS”, or in case of no previous BS,

“interested in BS” or “not interested in BS”. Other frequently used demographic parameters (age, sex, ethnic origin) were not included in the explorative models, since they were equally distributed among subgroups and >93% of all respondents were adult Caucasian females. Nagelkerke's R^2 was used to express the coefficient of determination of the models.

Results

Respondents

Out of 2387 unique visits at the landing page, 1601 respondents completed the study (completion rate: 67.1%). The survey was mostly filled in via smartphones (82.5%), and took in average 6min20sec to complete. The final study cohort included 1482 respondents (Supplemental Table 1.).

Free word associations

The total number of unique associations was 1793. After the spellcheck, lemmatization and merging, the number of associations decreased to 235, out of which 84 were non-idiosyncratic and represented the basis of the network analysis. The co-occurring network of associations is presented in Fig. 1. Fig 2. presents the respective emotional labels, whereas respondents' characteristics are shown in Table 1. Postbariatric patients had a significantly higher proportion of positive emotional labels than non-operated adults with obesity (79% vs 75%, $t(1480)=1.82$; $P = 0.002$, $d=0.18$). We labeled the two cognitive modules as benefit- or procedure-focused, based on their most frequent associations. Significant attractive and repulsive connections between associations are presented visually in Supplemental Fig. 2. The associations with the lowest community attachment (measured by local modularity) were termed “bridging” associations, since they were frequently mentioned by

respondents of both groups (Supplemental Fig. 3). These included for example: “slim”, “excess skin” and “pride”. The association “new life” (most frequent emotional labels: joy, gratitude, hope, pride and surprise) resulted in a stand-alone module, which might represent the ultimate bridging association between the two cognitive modules.

Prediction models of BMI status, BS status and interest in BS

Results of the regression models are presented in Table 2. Respondents with a current \geq Class II obesity were more likely to have a benefit-focused perception of weight loss, to mention hope more frequently, and gratitude less frequently than respondents without severe obesity. Respondents who had undergone BS were more likely to belong to the procedure-focused cognitive module and to mention pride, surprise and especially gratitude more frequently than respondents without history of BS. In the subgroup of respondents without previous BS, those who were interested in undergoing BS were more likely to express a benefit-focused mindset and use more frequently *hope* as emotional label.

Body Image Dissatisfaction

The BID score is presented according to the cognitive module membership in Table 1. Overall, current BMI showed a) a strong correlation with BID (Pearson's $r = 0.6$, $P < 0.001$, Supplemental Fig. 4), b) an even stronger correlation with current body image ($r = 0.77$, $P < 0.001$) and c) a weaker correlation with the expected body image ($r = 0.42$, $P < 0.001$). ~~whereas~~ Sex-related differences were not identified (BID = 2.3 ± 1.7 for females and 2.3 ± 1.7 for males). Only 10.9% of all respondents ($n = 162$) were free of BID (score = 0).

Respondents who would not consider to undergo BS

Out of the 409 respondents without prior BS, 66 stated to have no interest to undergo BS. This subgroup had a mean age of 43.6 years and consisted of Caucasian (91%) females (92%) with a BID of 2.2 who managed to lose weight non-surgically (highest lifetime BMI: $45.4 \pm 8.3 \text{ kg/m}^2$, current BMI: $37.9 \pm 8.8 \text{ kg/m}^2$). Only 53% of this group had regular contact with BS patients.

Discussion

This study achieved data-driven interpretation of the perception of weight loss in a large cohort of adults, mainly Caucasian women, with obesity and/or history of BS. This is the first application of this novel, network-analysis based methodology in the context of BS, where patients' mindset and emotional demeanor play a cardinal role in therapeutic decision-making.

A major finding was that the perception of "weight loss" in adults with obesity formed two distinct mental representations or modules. Some respondents predominantly mentioned associations related to the procedure of weight loss (i.e.: *sport, diet, surgery, effort* and *sacrifice*), while others provided associations related to the benefits of weight loss (i.e.: *health, activity, agility, attractiveness, happiness* and *quality or joy of life*). In analogy to the latent class analysis of a recent study on patient preferences for BS [22], the larger module was labelled as *benefit-focused*, and the smaller as *procedure-focused*. The above-mentioned American study identified an additional, "cost-sensitive" mindset among BS candidates, however, in the current study, the financial burdens of BS were not mentioned. This is most likely explained by the inherent differences between the health-

215 insurance systems of German-speaking European countries and the complex structure of health-
insurance coverage in the United States of America, where private funding is often involved. The most
frequently mentioned association was “*health*”, confirming from the patients’ perspective the utmost
importance of the metabolic effects of weight loss. Several associations highlighted the social
pressure related to weight loss (i.e.: *social life, employment, normality, friendship, confidence, new*
220 *life*). This observation complements the findings of a previous free-word association study in the
context of obesity, in which healthy students had to provide their impressions on *fatness*. Surprisingly,
they linked the fatness-related bad feelings with social pressure rather than with the health or self-
comfort of the people with obesity [23].

Explorative models were used to test whether cognitive module membership was associated
225 with previous or planned BS. Respondents with a history of BS adhered more frequently to the
procedure-focused cognitive module, mirroring their pragmatic approach to weight loss. In contrast,
those with an interest to undergo BS belonged more likely to the abstract, benefit-focused cognitive
module. It has been shown earlier that abstract (vs. concrete) representations of ideal end-states are
likely to influence engagement in a specific goal-oriented behavior [24]. In the context of obesity,
230 optimistic expectations were previously shown to favor higher rates of successful weight loss
following behavioral weight reduction [25]. However, unrealistic expectations can be responsible for
the cycle of ~~repeated failures~~sub-optimal outcomes and renewed efforts to achieve self-change (i.e.:
false-hope syndrome, “yo-yo effect”)[26]. Therefore, the decision of undergoing BS may reflect a new
approach to weight loss by patients who already experienced several unsuccessful conservative
235 attempts to tackle obesity [1]. Although body image is an important indicator of patients’ wellbeing
after BS [27], BID ratings did not differ between the two cognitive modules, but showed a very

significant correlation with current BMI, and were higher in participants who planned to undergo BS, compared to those who did not, or already had BS.

Based on our findings, the emotional component may play a more important role than the cognitive component in the decision-making process of obese patients who seek to engage in new health behaviors. In line with our hypothesis, history of BS favored the use of positive emotional labels, however, the effect size was rather small. In fact, all respondents selected mainly positive emotions to label their associations. Remarkably, patients willing to undergo BS were more likely to express *hope*, while patients after BS expressed *gratitude* and *pride* more frequently, ~~suggests~~ suggesting that BS may fundamentally change the individuals' affective connection with weight loss. The emotional state of hope is forward-looking and may characterize both the procedures and the benefits related to a planned action, reflected by its equal distribution between the two cognitive modules [28]. Respondents with a \geq Class II obesity mentioned *hope* more frequently, reflecting their unmet need of achieving weight loss. In contrast, *gratitude* and *pride* were more frequently mentioned in the benefit-focused module and by those who already had BS and had a current BMI <35 kg/m². The feeling of gratitude is one of the most typical responses to perceived benevolence, which appears to foster prosocial behaviors and correlates with other factors of psychological well-being and agreeableness [29]. Pride is a subjective status-related emotion, which has a fundamental affective role in status seeking, attainment, and signaling [30]. According to a Brazilian study in patients experiencing postbariatric weight regain, the feeling of gratitude toward BS remained, whereas pride was replaced by a feeling of ~~failure~~ sub-optimal outcomes and low self-esteem [31]. *Peacock et al.* also found that the extent of emotional component in bariatric candidates' motivation (by comparing desperate patients ("have to") with those who expressed sentiment of being tired

("don't want to") or of pragmatism ("want to")) was associated with better postbariatric weight loss [8]. Subsequently, postoperative weight loss was shown to have a strong positive impact on patients' psychological well-being (including: autonomy, environmental mastery, personal growth, positive relations with others, purpose in life, and self-acceptance), and even on marital satisfaction [32].

We acknowledge that associations are determined by a variety of personal experiences and future projections, and are not necessarily anchored to one single factor (such as BMI or BS).

Nevertheless, this study provided a comprehensive structure of ~~obese individuals'~~ adults' with obesity free expressions about weight loss, enhancing our understanding of patients' priorities and ~~may~~ contributeing to the improvement of health care-related communicational schemes. Discrepant expectations may significantly increase postoperative dissatisfaction with body image and decrease the extent of achievable weight loss [33], therefore setting realistic preoperative expectations and stepwise postoperative goals are crucial. Our findings also promote the collaborative approach between healthcare providers to support evidence-based treatments of obesity [34]. Focusing on the *consumer value* is the first principle of the *lean thinking* health care management method [35]. Association network models have been proven suitable for studying consumer-based brand equity by identifying relevant "character traits" which may assist the optimization of current processes [36]. In the context of BS, *Powell et al.* showed that psychological skills interventions can be implemented successfully to improve postbariatric patients' participation in exercise programs [37]. Consequently, the network-based map of associations to weight loss presented in this study may serve as a navigational tool for caregivers, policy actors and researchers to better identify patients' implicit interest toward weight loss and to design new interventions aiming to tackle mindset barriers in the acceptance of BS among patients with obesity [38]. Our findings suggest that complementing factual

information on BS with emotional experiences of postbariatric patients (i.e.: gratitude and pride) and emphasizing the benefits rather than the procedural aspects of achieving weight loss may be effective in changing perspectives on life-priorities of people with obesity. As there is a growing interest for health-related information on multiple digital platforms, including social media, the relevant positioning of “weight loss” can influence large groups of people with obesity and change social norms [39]. Once a mindset has been primed or activated, there is an increased likelihood that it will be used in upcoming tasks to interpret new information and to define subsequent actions [24]. A subgroup of the respondents stated not to be interested to undergo BS, nevertheless, by following BS-related social media they manifested interest towards BS and seem to be seeking facts and/or patient narratives to assist their health-related decisions.

The main limitation of the study derives from the inherent methodologic bias of internet-based surveys: 1) the non-representative nature of the Internet population and 2) the self-selection of participants (volunteer effect) [15]. These factors may explain the very high proportion of Caucasian females with previous BS within all respondents. In the absence of longitudinal data collection, demographic parameters – especially current BMI – are likely to confound the relationship of BS and cognitive module membership. Although there are techniques to account for the effect of BMI (i.e.: propensity-score matching), we aimed to demonstrate real-world scenarios where the stage of obesity and BS history are strongly interconnected. The lack of directly measured BMI values represent an additional imperfection. Nevertheless, self-reported height and weight were found to highly correlate with measured values ($R \geq 0.9$) in the Danish Health Survey [40].

Conclusion

The combination of research methods (data-driven cross-network analysis of free associations with emotional valence and use of validated psychologic questionnaire) was synergistic in mapping patients' motivational drivers, beliefs, attitudes or feelings towards weight loss. Respondents with severe obesity and those planning to undergo BS are more likely to be benefit-focused and to express "hope", compared to their slimmer postbariatric peers, who are more often procedure-focused, feel "gratitude" and "pride" more often and suffer less from body image dissatisfaction. Emotional and mental schemes stemming from patients' free associations may enrich the informed consent process prior to BS and could serve as basis for the development of novel communicational strategies between health-care providers and patients with obesity.

Acknowledgments

We are indebted to **Blinded**, PhD candidate at the Doctoral School of Psychology, **Blinded**, for her advices regarding the interpretation of the results. Further, we would like to thank the help of **Blinded**, administrator of the Adipositas Zürich Facebook group, for his contribution in disseminating the invitations for the survey. We are also grateful to all respondents for their time and valuable input. This work was supported by the assistant-professorship research grant awarded by the **Blinded**; **Blinded**, MSc was supported by **Blinded** New National Excellence Program of the **Blinded**. The sponsors had no influence on the study design, or collection, analysis and interpretation of data.

Disclosure: The authors declare no conflict of interest. Prof. **Blinded** reports personal fees from Johnson & Johnson and Medtronic, outside the submitted work.

Ethical approval

325 The Cantonal Ethics Committee of Zurich stated that the study did not fall under the scope of the Swiss Human Research Act, and consequently, does not need ethical approval (BASEC-Nr. Req-2017-00903).

Tables

330 **Table 1.** Respondents' characteristics according to cognitive module membership. *BMI: Body Mass Index. BS: bariatric surgery. *: significant ($P < 0.01$) difference in the frequency of the emotion between the two modules.*

335 **Table 2.** Explorative models using logistic regression investigating the effect of the stage of obesity and of bariatric surgery status on cognitive module membership and use of different emotional labels. *CI: confidence interval, BMI: body mass index, BS: bariatric surgery*

A. Respondents with \geq WHO Class II Obesity vs. Respondents with $<$ WHO Class II Obesity (= reference). Based on the entire cohort [$n = 1482$; $R^2 = 0.156$]

340 B. Respondents who underwent bariatric surgery vs. Respondents without history of bariatric surgery (= reference). Based on the entire cohort [$n = 1482$, $R^2 = 0.169$]

345 C. Respondents with an interest to undergo bariatric surgery vs. Respondents not interested to undergo bariatric surgery (= reference). Within the subgroup of respondents without history of bariatric surgery [$n = 409$; $R^2 = 0.142$]

Figures

350 **Figure 1.** Free-word association networks to “weight loss” in patients with obesity. The modules reflect the data-driven frequency of co-occurrences of associations. Potential interpretation: *Green: benefit-focused mindset; Purple: procedure-focused mindset.*

Figure 2. Frequency of emotional labels assigned to the associations of each cognitive module

355

Supplemental Material

Supplemental Methods

Supplemental Method 1. The survey as it was presented to the participants (MethodS1.pdf)

Supplemental Table

Supplemental Table 1.

A. Descriptive characteristics of the respondents based on their current stage of obesity and on their bariatric surgery status. BMI: body mass index

B. Comparison of demographic parameters between subgroups

~~Descriptive characteristics of the respondents based on their current stage of obesity and on their bariatric surgery status. BMI: body mass index~~

Supplemental Figures

Supplemental Figure 1. Flowchart of the selection of the study cohort

Supplemental Figure 2. Attractive and repulsive associations

Supplemental Figure 3. Bridging associations

Supplemental Figure 4. Correlation of current BMI and Body Image Dissatisfaction

References

- [1] Bray GA, Fruhbeck G, Ryan DH, Wilding JP. Management of obesity. *Lancet*. 2016;387:1947-56.
- 375 [2] Kaplan LM, Golden A, Jinnett K, Kolotkin RL, Kyle TK, Look M, et al. Perceptions of Barriers to Effective Obesity Care: Results from the National ACTION Study. *Obesity (Silver Spring)*. 2018;26:61-9.
- [3] Puhl RM, Heuer CA. Obesity stigma: important considerations for public health. *Am J Public Health*. 2010;100:1019-28.
- [4] Gero D, Hulesch B, Bueter M. Mindset and Communication Barriers in the Diffusion of Bariatric Surgery. *Curr Atheroscler Rep*. 2018;20:38.
- 380 [5] Munoz DJ, Lal M, Chen EY, Mansour M, Fischer S, Roehrig M, et al. Why patients seek bariatric surgery: a qualitative and quantitative analysis of patient motivation. *Obes Surg*. 2007;17:1487-91.
- [6] Libeton M, Dixon JB, Laurie C, O'Brien PE. Patient motivation for bariatric surgery: characteristics and impact on outcomes. *Obes Surg*. 2004;14:392-8.
- 385 [7] Edward KL, Hii MW, Giandinoto JA, Hennessy J, Thompson L. Personal Descriptions of Life Before and After Bariatric Surgery From Overweight or Obese Men. *Am J Mens Health*. 2018;12:265-73.
- [8] Peacock JC, Perry L, Morien K. Bariatric patients' reported motivations for surgery and their relationship to weight status and health. *Surg Obes Relat Dis*. 2018;14:39-45.
- [9] Sharman MJ, Venn AJ, Hensher M, Wilkinson S, Palmer AJ, Williams D, et al. Motivations for Seeking
- 390 Bariatric Surgery: The Importance of Health Professionals and Social Networks. *Bariatr Surg Pract P*. 2016;11:104-9.
- [10] Walach H, Loughlin M. Patients and agents - or why we need a different narrative: a philosophical analysis. *Philos Ethics Humanit Med*. 2018;13:13.
- [11] Teutsch C. Patient-doctor communication. *Med Clin North Am*. 2003;87:1115-45.
- 395 [12] File B, Keczer Z, Vancso A, Bothe B, Toth-Kiraly I, Hunyadi M, et al. Emergence of polarized opinions from free association networks. *Behav Res Methods*. 2018.
- [13] Luduena GA, Behzad MD, Gros C. Exploration in free word association networks: models and experiment. *Cogn Process*. 2014;15:195-200.
- [14] Vijayasarathi A, Kharkar R, Salamon N. Strategies for Patient-Centered Communication in the Digital Age.
- 400 *Curr Probl Diagn Radiol*. 2018.
- [15] Eysenbach G. Improving the quality of Web surveys: the Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res*. 2004;6:e34.
- [16] Ekman P. Are there basic emotions? *Psychol Rev*. 1992;99:550-3.
- [17] Robinson DL. Brain function, emotional experience and personality. *Netherlands Journal of Psychology*.
- 405 2008;64:152-68.
- [18] Blanch L. Under Construction: Body Image and Self-Representation in Virtual Worlds. . Toronto, Canada: York University, Toronto, Canada; 2014.
- [19] Ratcliff MB, Eshleman KE, Reiter-Purtill J, Zeller MH. Prospective changes in body image dissatisfaction among adolescent bariatric patients: the importance of body size estimation. *Surg Obes Relat Dis*.
- 410 2012;8:470-5.
- [20] Research AAFPO. Standard Definitions: Final Dispositions of Case Codes and Outcome Rates for Surveys. 8th ed. Oakbrook Terrace, IL, USA2015.
- [21] Lancichinetti A, Fortunato S. Consensus clustering in complex networks. *Sci Rep*. 2012;2:336.
- [22] Rozier MD, Ghaferi AA, Rose A, Simon NJ, Birkmeyer N, Prosser LA. Patient Preferences for Bariatric
- 415 Surgery: Findings From a Survey Using Discrete Choice Experiment Methodology. *JAMA Surg*. 2018:e184375.
- [23] Mondragon NI, Txertudi MB. Understanding fatness in the public sphere of young students: social representations and emotional response. *Cad Saude Publica*. 2018;34:e00197917.
- [24] Torelli CJ, Kaikati AM. Values as predictors of judgments and behaviors: the role of abstract and concrete mindsets. *J Pers Soc Psychol*. 2009;96:231-47.

420 [25] Oettingen G, Wadden TAJCT, Research. Expectation, fantasy, and weight loss: Is the impact of positive thinking always positive? 1991;15:167-75.

[26] Trottier K, Polivy J, Herman CP. Effects of resolving to change one's own behavior: expectations vs. experience. *Behav Ther.* 2009;40:164-70.

425 [27] Monpellier VM, Antoniou EE, Mulkens S, Janssen IMC, van der Molen ABM, Jansen ATM. Body image dissatisfaction and depression in postbariatric patients is associated with less weight loss and a desire for body contouring surgery. *Surg Obes Relat Dis.* 2018;14:1507-15.

[28] Gallagher MWCJ. *The Oxford handbook of hope.* New York, NY: Oxford University Press; 2018.

[29] McCullough ME, Kilpatrick SD, Emmons RA, Larson DB. Is gratitude a moral affect? *Psychol Bull.* 2001;127:249-66.

430 [30] Bollo H, Bothe B, Toth-Kiraly I, Orosz G. Pride and Social Status. *Front Psychol.* 2018;9:1979.

[31] Carvalho A, Jr., Turato ER, Chaim EA, Magdaleno R, Jr. Weight regain among women after metabolic and bariatric surgery: a qualitative study in Brazil. *Trends Psychiatry Psychother.* 2014;36:140-6.

[32] Karagulle OO, Yavuz E, Gulcicek OB, Solmaz A, Senturk S, Erdogan A, et al. Psychological well-being and marital satisfaction in response to weight loss after bariatric surgery. *Surg Today.* 2019.

435 [33] Conceição EM, de Lourdes M, Moreira L, Pinto-Bastos A, Félix S. Weight loss expectations and weight loss after surgery: the mediating role of body image and weight concerns. *Surgery for Obesity and Related Diseases.* 2020.

[34] Sharma AM, Belanger A, Carson V, Krah J, Langlois MF, Lawlor D, et al. Perceptions of barriers to effective obesity management in Canada: Results from the ACTION study. *Clin Obes.* 2019;9:e12329.

440 [35] Aij KH, Teunissen M. Lean leadership attributes: a systematic review of the literature. *J Health Organ Manag.* 2017;31:713-29.

[36] Teichert TA, Schöntag K. Exploring consumer knowledge structures using associative network analysis. 2010;27:369-98.

[37] Powell SM, Fasczewski KS, Gill DL, Davis PG. Go with the FLOW: Implementation of a psychological skills intervention in an exercise program for post-bariatric surgery patients. *J Health Psychol.* 2018;1359105318793182.

445 [38] Nowicki T, Burns C, Fulbrook P, Jones J. Changing the mindset: an inter-disciplinary approach to management of the bariatric patient. *Collegian.* 2009;16:171-5.

[39] Vallone D, Greenberg M, Xiao H, Bennett M, Cantrell J, Rath J, et al. The Effect of Branding to Promote Healthy Behavior: Reducing Tobacco Use among Youth and Young Adults. *Int J Environ Res Public Health.* 2017;14.

450 [40] Neermark S, Holst C, Bisgaard T, Bay-Nielsen M, Becker U, Tolstrup JS. Validation and calibration of self-reported height and weight in the Danish Health Examination Survey. *Eur J Public Health.* 2018.

Title

Mental and emotional representations of “weight loss”: free-word association networks in members of bariatric surgery-related social media communities

5 Authors

Blinded

Manuscript category: Integrated Health Article

10 **Communication:** The study was presented in part at the 24th International Federation for the Surgery of Obesity and Metabolic Disorders World Congress on 5 September 2019 in Madrid, Spain and at the 106th Annual meeting of the Swiss Society of Surgeons on 17 May 2019 in Bern, Switzerland.

Funding: This work was supported by the assistant-professorship research grant awarded by the **Blinded** to Prof. Dr. med. **Blinded**, PhD; **Blinded**, MSc was supported by the **Blinded** New
15 National Excellence Program of the **Blinded**.

Correspondence to:

Blinded

Running title: Mental and emotional representations of weight loss

20 **Abstract**

Background: Mindset and communication barriers may hinder the acceptance of bariatric surgery (BS) by the eligible patient population.

Objectives: To improve the understanding of expectations, opinions, emotions and attitudes toward weight-loss among patients with obesity.

25 **Setting:** Switzerland, Germany, Austria.

Methods: Survey data collected from BS-related social media communities (N=1482). Participants were asked to write 5 words that first came to their mind about “weight loss”, and to select 2 emotions which best described their corresponding feelings. Demographic and obesity-related data were collected. Cognitive representations were constructed based on the co-occurrence network of
30 associations, using validated data-driven methodology.

Results: Respondents were Caucasian (98%), female (94%), aged 42.5 ± 10.1 years, current/highest lifetime body mass index = $36.9 \pm 9 / 50.7 \pm 8.7$ kg/m². The association network analysis revealed two cognitive modules: benefit-focused (health, attractiveness, happiness, agility) and procedure-focused (effort, diet, sport, surgery). Patients willing to undergo BS were more benefit-focused (Odds ratio (OR)=2.4, P=0.02) and expressed more ‘hope’ (OR=142, P<0.001). History of BS was associated with
35 higher adherence to the procedure-focused module (OR=2.3, P<0.001), and with increased use of the emotions ‘gratitude’ (OR=107, P<0.001), ‘pride’ (OR=15, P<0.001), and decreased mention of ‘hope’ (OR=0.03, P<0.001).

Conclusion: Patients with obesity in our study tend to think about weight loss along two cognitive
40 schemes, either emphasizing its expected benefits or focusing on the process of achieving it. Benefit-focused respondents were more likely to consider BS, and to express hope rather than gratitude or pride. Novel communication strategies may increase the acceptance of BS by incorporating weight loss-related cognitive and emotional content stemming from patients’ free associations.

Keywords

45 Bariatric Surgery; Weight loss; Free-word Association; Network; Obesity; Emotional valence; Mindset; Motivation; Body image dissatisfaction

Introduction

Bariatric surgery (BS) is currently the most effective modality in the treatment of severe obesity and related diseases [1]. Despite its growing worldwide popularity as a treatment option, BS is rather underused by the eligible patient population [2]. Mindset and communication barriers seem to influence the diffusion of BS: lack of information, incorrect information, and stigmatization of individuals with obesity have proven to contribute to sub-optimal use of health care by the eligible patients and to insufficient cooperation between primary care physicians and BS centers [3, 4]. This may also hinder the identification and referral of eligible patients and undermine optimal postbariatric follow-up [4].

Available literature on motivational drivers in favor of seeking BS remains scarce and stems mainly from small cohorts [5-7]. Nevertheless, corroborative evidence shows that the willingness to improve health is the cornerstone for seeking BS, contextualized with psychological, emotional, environmental and social factors [8, 9]. Within the developing framework of person-centered care, the patients' narrative is a key factor in improving decision-making [10]. However, due to the sensitive, personal and private nature of patients' perception of their own body weight, direct questions from healthcare professional may be perceived as judgmental or scolding, and may rapidly close down communication [11]. In this sensitive context, an anonymized online survey with a free word association task seems to be an optimal approach to scope group opinion [12].

Free word association is a widely used technique in market research and psychology to encourage respondents to express openly their underlying motivations, beliefs, attitudes or feelings regarding a specific topic [13]. This technique enhances the unconstrained expressions of respondents and overcomes limitations of predefined questionnaires.

We aimed to map the mental and emotional representation of the term *weight loss* in patients with obesity who had a documented interest in BS, using free word associations combined with a novel data-driven associative analytical method [12]. Further, we aimed to investigate the relationship between these representations and clinical parameters, such as stage of obesity and previous or planned experience with BS. We hypothesized that BS had a “positive” impact on the emotional labeling of *weight loss* related free word associations. Our findings may serve as basis to improve patient-centered communication strategies, online and in clinical practice alike [14].

Methods

We conducted an anonymized online survey to understand how adults with severe obesity (those who seek BS, postbariatric patients, and those who are not interested to undergo BS) perceive weight loss. To ensure that participants are familiar with BS, respondents were recruited on a voluntary basis from bariatric surgery-related German-speaking social media groups between 02/11/2018 and 04/04/2018. An invitation for participation and three reminders every two weeks were posted on the ‘Adipositas Zürich’ (4100 followers) and ‘Adipositas Chirurgie - Fragen und Antworten’ (1500 followers) Facebook© groups, and respondents were encouraged to share the invitation within their own social networks. Thus, the survey was “open” to any respondents with a presumed interest in BS. We aimed to enroll a convenience sample of >500 respondents. The survey was administered via Typeform™ (Barcelona, Spain) (Supplemental Methods 1.), and only complete questionnaires could be submitted. Respondents could not be backtracked to a specific Facebook© profile. The Cantonal Ethics Committee of Zurich concluded that this study did not fall under the scope of the Swiss Human Research Act, thus ethical approval was not required. The manuscript was

prepared according to the Checklist for Reporting Results of Internet E-Surveys (CHERRIES) guidelines [15]. Network analyses were performed using Matlab 2017 (The Mathworks); logistic regressions and correlations were performed using R Software for Statistical Computing v3.5.2 (R Foundation, Vienna, Austria).

95 Building the dataset

Basic Demographics

Respondents were asked to self-report their age, sex, ethnicity, civil status, height, maximal and current body weight, past or planned bariatric operations, previous non-surgical weight loss attempts and their social interactions with bariatric patients. Body mass index (BMI) was calculated
100 by the investigators based on the anthropometric data provided by the participants.

Free word associations

Participants were asked to write five words, which first came to their mind about “weight loss”. Further, they had to select two emotions from a list of 20 basic emotions (interest, anxiety, empathy, contempt, surprise, indifference, hope, fear, gratitude, anger, joy, sadness, calmness,
105 frustration, pride, shame, generosity, envy, sympathy, antipathy) that best described their feelings about each of the five associations they provided [16, 17]. Emotional labeling was successfully applied in a recent study to improve the interpretation of free-word associations by the addition of affective content [12].

Body image dissatisfaction (BID)

110 The color version of the reliable and validated Stunkard Figure Rating Scale was used to assess current and ideal body size estimations [18]. This instrument uses a series of 9 gender-specific

silhouettes that gradually increase in body size. Although developed for normal weight adults with traditional eating disorders (e.g.: anorexia nervosa), this scale has been previously applied in the context of BS [19]. Respondents were instructed to select the figure that indicates (a) how they look currently and (b) how they would like to look like. A discrepancy score indicating *Body Image Dissatisfaction* (BID) was calculated by subtracting scores of the ideal figure (b) from the current (a) figure. Lower discrepancy scores were interpreted as lower body image dissatisfaction.

Data pre-processing

Response rate was calculated based on definitions and metrics proposed by the American Association for Public Opinion Research [20]. Respondents who provided incoherent data (i.e.: highest lifetime BMI < current BMI) or were found to be outside the target group (i.e.: highest lifetime BMI < 35 kg/m², history of gastric banding, etc.) were excluded from the study cohort in a step-wise manner (Supplemental Fig. 1). Free word associations to “weight loss” were first spellchecked and lemmatized (i.e.: to return the base of a word) by two independent native German speakers. Associations were merged if their English translation was identical. Idiosyncratic expressions (associations with <9 occurrences) were left out.

Association network analysis

In order to extract the most prominent mental representations from the numerous individual associations provided by the respondents, we applied an in-house developed algorithm, using a network-based methodology validated and described earlier by our group [12]. To create such a network, nodes (free word associations) and edges (statistical co-occurrences between two associations) were determined. Statistical co-occurrence was calculated by log-likelihood ratios (LLR).

The LLR value can denote attractive (when two associations are often mentioned together) and repulsive connections (when two associations are rarely mentioned together) with a corresponding level of significance. In the network of statistically related associations, modularity maximization based consensus clustering algorithm was applied to explore any densely connected subnetworks (i.e.: modules) [21]. We have previously demonstrated that these modules of associations - as consistent patterns in individual association sequences – are able to capture the most prominent cognitive representation of a given cue [12]. Associations with weak module attachment were separately identified, since they may serve as bridges in future communicational strategies to simultaneously address individuals with different cognitive modular membership.

Qualitative and quantitative analysis of the map

First, we used descriptive statistics to characterize the subgroups of interests, as well as the members of the two cognitive modules. *Second*, we created explorative models by logistic regression to describe the relationship between free-word associations (cognitive module membership, emotional labels) and clinical parameters (stage of obesity, BS status, interest to undergo BS). The cognitive module of respondents was determined by the association cluster to which the majority of their associations belonged. Emotions were expressed as frequency (%) and the unfrequently (<4%) mentioned emotions were not included in the models to decrease false discovery rate. Respondents' current BMI status was dichotomized according to World Health Organisation obesity class II (\geq / $<$ 35 kg/m²), since this is the currently recommended cutoff for being eligible to BS [1]. We also created subgroups based on the BS status of respondents: “underwent BS”, or in case of no previous BS, “interested in BS” or “not interested in BS”. Other frequently used demographic parameters (age, sex, ethnic origin) were not included in the explorative models, since they were equally distributed among

subgroups and >93% of all respondents were adult Caucasian females. Nagelkerke's R^2 was used to express the coefficient of determination of the models.

Results

Respondents

Out of 2387 unique visits at the landing page, 1601 respondents completed the study (completion rate: 67.1%). The survey was mostly filled in via smartphones (82.5%), and took in average 6min20sec to complete. The final study cohort included 1482 respondents (Supplemental Table 1.).

Free word associations

The total number of unique associations was 1793. After the spellcheck, lemmatization and merging, the number of associations decreased to 235, out of which 84 were non-idiosyncratic and represented the basis of the network analysis. The co-occurring network of associations is presented in Fig. 1. Fig 2. presents the respective emotional labels, whereas respondents' characteristics are shown in Table 1. Postbariatric patients had a significantly higher proportion of positive emotional labels than non-operated adults with obesity (79% vs 75%, $t(1480)=1.82$; $P = 0.002$, $d=0.18$). We labeled the two cognitive modules as benefit- or procedure-focused, based on their most frequent associations. Significant attractive and repulsive connections between associations are presented visually in Supplemental Fig. 2. The associations with the lowest community attachment (measured by local modularity) were termed "bridging" associations, since they were frequently mentioned by respondents of both groups (Supplemental Fig. 3). These included for example: "slim", "excess skin" and "pride". The association "new life" (most frequent emotional labels: joy, gratitude, hope, pride

and surprise) resulted in a stand-alone module, which might represent the ultimate bridging association between the two cognitive modules.

Prediction models of BMI status, BS status and interest in BS

Results of the regression models are presented in Table 2. Respondents with a current \geq Class II obesity were more likely to have a benefit-focused perception of weight loss, to mention hope more frequently, and gratitude less frequently than respondents without severe obesity. Respondents who had undergone BS were more likely to belong to the procedure-focused cognitive module and to mention pride, surprise and especially gratitude more frequently than respondents without history of BS. In the subgroup of respondents without previous BS, those who were interested in undergoing BS were more likely to express a benefit-focused mindset and use more frequently *hope* as emotional label.

Body Image Dissatisfaction

The BID score is presented according to the cognitive module membership in Table 1. Overall, current BMI showed a) a strong correlation with BID (Pearson's $r = 0.6$, $P < 0.001$, Supplemental Fig. 4), b) an even stronger correlation with current body image ($r = 0.77$, $P < 0.001$) and c) a weaker correlation with the expected body image ($r = 0.42$, $P < 0.001$). Sex-related differences were not identified (BID = 2.3 ± 1.7 for females and 2.3 ± 1.7 for males). Only 10.9% of all respondents ($n = 162$) were free of BID (score = 0).

Respondents who would not consider to undergo BS

Out of the 409 respondents without prior BS, 66 stated to have no interest to undergo BS. This subgroup had a mean age of 43.6 years and consisted of Caucasian (91%) females (92%) with a BID of 2.2 who managed to lose weight non-surgically (highest lifetime BMI: $45.4 \pm 8.3 \text{ kg/m}^2$, current BMI: $37.9 \pm 8.8 \text{ kg/m}^2$). Only 53% of this group had regular contact with BS patients.

Discussion

This study achieved data-driven interpretation of the perception of weight loss in a large cohort of adults, mainly Caucasian women, with obesity and/or history of BS. This is the first application of this novel, network-analysis based methodology in the context of BS, where patients' mindset and emotional demeanor play a cardinal role in therapeutic decision-making.

A major finding was that the perception of "weight loss" in adults with obesity formed two distinct mental representations or modules. Some respondents predominantly mentioned associations related to the procedure of weight loss (i.e.: *sport, diet, surgery, effort* and *sacrifice*), while others provided associations related to the benefits of weight loss (i.e.: *health, activity, agility, attractiveness, happiness* and *quality or joy of life*). In analogy to the latent class analysis of a recent study on patient preferences for BS [22], the larger module was labelled as *benefit-focused*, and the smaller as *procedure-focused*. The above-mentioned American study identified an additional, "cost-sensitive" mindset among BS candidates, however, in the current study, the financial burdens of BS were not mentioned. This is most likely explained by the inherent differences between the health-

insurance systems of German-speaking European countries and the complex structure of health-insurance coverage in the United States of America, where private funding is often involved. The most frequently mentioned association was “*health*”, confirming from the patients’ perspective the utmost importance of the metabolic effects of weight loss. Several associations highlighted the social pressure related to weight loss (i.e.: *social life, employment, normality, friendship, confidence, new life*). This observation complements the findings of a previous free-word association study in the context of obesity, in which healthy students had to provide their impressions on *fatness*. Surprisingly, they linked the fatness-related bad feelings with social pressure rather than with the health or self-comfort of the people with obesity [23].

Explorative models were used to test whether cognitive module membership was associated with previous or planned BS. Respondents with a history of BS adhered more frequently to the procedure-focused cognitive module, mirroring their pragmatic approach to weight loss. In contrast, those with an interest to undergo BS belonged more likely to the abstract, benefit-focused cognitive module. It has been shown earlier that abstract (vs. concrete) representations of ideal end-states are likely to influence engagement in a specific goal-oriented behavior [24]. In the context of obesity, optimistic expectations were previously shown to favor higher rates of successful weight loss following behavioral weight reduction [25]. However, unrealistic expectations can be responsible for the cycle of sub-optimal outcomes and renewed efforts to achieve self-change (i.e.: false-hope syndrome, “yo-yo effect”)[26]. Therefore, the decision of undergoing BS may reflect a new approach to weight loss by patients who already experienced several unsuccessful conservative attempts to tackle obesity [1]. Although body image is an important indicator of patients’ wellbeing after BS [27], BID ratings did not differ between the two cognitive modules, but showed a very

significant correlation with current BMI, and were higher in participants who planned to undergo BS, compared to those who did not, or already had BS.

240 Based on our findings, the emotional component may play a more important role than the cognitive component in the decision-making process of obese patients who seek to engage in new health behaviors. In line with our hypothesis, history of BS favored the use of positive emotional labels, however, the effect size was rather small. In fact, all respondents selected mainly positive emotions to label their associations. Remarkably, patients willing to undergo BS were more likely to
245 express *hope*, while patients after BS expressed *gratitude* and *pride* more frequently, suggesting that BS may fundamentally change the individuals' affective connection with weight loss. The emotional state of hope is forward-looking and may characterize both the procedures and the benefits related to a planned action, reflected by its equal distribution between the two cognitive modules [28]. Respondents with a \geq Class II obesity mentioned *hope* more frequently, reflecting their unmet need of
250 achieving weight loss. In contrast, *gratitude* and *pride* were more frequently mentioned in the benefit-focused module and by those who already had BS and had a current BMI <35 kg/m². The feeling of gratitude is one of the most typical responses to perceived benevolence, which appears to foster prosocial behaviors and correlates with other factors of psychological well-being and agreeableness [29]. Pride is a subjective status-related emotion, which has a fundamental affective role in status
255 seeking, attainment, and signaling [30]. According to a Brazilian study in patients experiencing postbariatric weight regain, the feeling of gratitude toward BS remained, whereas pride was replaced by a feeling of sub-optimal outcomes and low self-esteem [31]. *Peacock et al.* also found that the extent of emotional component in bariatric candidates' motivation (by comparing desperate patients ("have to") with those who expressed sentiment of being tired ("don't want to") or of pragmatism

260 (“want to”)) was associated with better postbariatric weight loss [8]. Subsequently, postoperative weight loss was shown to have a strong positive impact on patients’ psychological well-being (including: autonomy, environmental mastery, personal growth, positive relations with others, purpose in life, and self-acceptance), and even on marital satisfaction [32].

We acknowledge that associations are determined by a variety of personal experiences and
265 future projections, and are not necessarily anchored to one single factor (such as BMI or BS). Nevertheless, this study provided a comprehensive structure of adults’ with obesity free expressions about weight loss, enhancing our understanding of patients’ priorities and contributing to the improvement of health care-related communicational schemes. Discrepant expectations may significantly increase postoperative dissatisfaction with body image and decrease the extent of
270 achievable weight loss [33], therefore setting realistic preoperative expectations and stepwise postoperative goals are crucial. Our findings also promote the collaborative approach between healthcare providers to support evidence-based treatments of obesity [34]. Focusing on the *consumer value* is the first principle of the *lean thinking* health care management method [35]. Association network models have been proven suitable for studying consumer-based brand equity by identifying
275 relevant “character traits” which may assist the optimization of current processes [36]. In the context of BS, *Powell et al.* showed that psychological skills interventions can be implemented successfully to improve postbariatric patients’ participation in exercise programs [37]. Consequently, the network-based map of associations to weight loss presented in this study may serve as a navigational tool for caregivers, policy actors and researchers to better identify patients’ implicit interest toward weight
280 loss and to design new interventions aiming to tackle mindset barriers in the acceptance of BS among patients with obesity [38]. Our findings suggest that complementing factual information on BS with

emotional experiences of postbariatric patients (i.e.: gratitude and pride) and emphasizing the benefits rather than the procedural aspects of achieving weight loss may be effective in changing perspectives on life-priorities of people with obesity. As there is a growing interest for health-related information on multiple digital platforms, including social media, the relevant positioning of “weight loss” can influence large groups of people with obesity and change social norms [39]. Once a mindset has been primed or activated, there is an increased likelihood that it will be used in upcoming tasks to interpret new information and to define subsequent actions [24]. A subgroup of the respondents stated not to be interested to undergo BS, nevertheless, by following BS-related social media they manifested interest towards BS and seem to be seeking facts and/or patient narratives to assist their health-related decisions.

The main limitation of the study derives from the inherent methodologic bias of internet-based surveys: 1) the non-representative nature of the Internet population and 2) the self-selection of participants (volunteer effect) [15]. These factors may explain the very high proportion of Caucasian females with previous BS within all respondents. In the absence of longitudinal data collection, demographic parameters – especially current BMI – are likely to confound the relationship of BS and cognitive module membership. Although there are techniques to account for the effect of BMI (i.e.: propensity-score matching), we aimed to demonstrate real-world scenarios where the stage of obesity and BS history are strongly interconnected. The lack of directly measured BMI values represent an additional imperfection. Nevertheless, self-reported height and weight were found to highly correlate with measured values ($R \geq 0.9$) in the Danish Health Survey [40].

Conclusion

The combination of research methods (data-driven cross-network analysis of free associations
305 with emotional valence and use of validated psychologic questionnaire) was synergistic in mapping
patients' motivational drivers, beliefs, attitudes or feelings towards weight loss. Respondents with
severe obesity and those planning to undergo BS are more likely to be benefit-focused and to express
"hope", compared to their slimmer postbariatric peers, who are more often procedure-focused, feel
"gratitude" and "pride" more often and suffer less from body image dissatisfaction. Emotional and
310 mental schemes stemming from patients' free associations may enrich the informed consent process
prior to BS and could serve as basis for the development of novel communicational strategies between
health-care providers and patients with obesity.

Acknowledgments

315 We are indebted to **Blinded**, PhD candidate at the Doctoral School of Psychology, **Blinded**, for her
advices regarding the interpretation of the results. Further, we would like to thank the help of
Blinded, administrator of the Adipositas Zürich Facebook group, for his contribution in
disseminating the invitations for the survey. We are also grateful to all respondents for their time and
valuable input. This work was supported by the assistant-professorship research grant awarded by
320 the **Blinded**; **Blinded**, MSc was supported by **Blinded** New National Excellence Program of the
Blinded. The sponsors had no influence on the study design, or collection, analysis and
interpretation of data.

Disclosure: The authors declare no conflict of interest. Prof. **Blinded** reports personal fees from Johnson & Johnson and Medtronic, outside the submitted work.

325 **Ethical approval**

The Cantonal Ethics Committee of Zurich stated that the study did not fall under the scope of the Swiss Human Research Act, and consequently, does not need ethical approval (BASEC-Nr. Req-2017-00903).

Tables

330 **Table 1.** Respondents' characteristics according to cognitive module membership. *BMI: Body Mass Index. BS: bariatric surgery. *: significant ($P < 0.01$) difference in the frequency of the emotion between the two modules.*

335 **Table 2.** Explorative models using logistic regression investigating the effect of the stage of obesity and of bariatric surgery status on cognitive module membership and use of different emotional labels. *CI: confidence interval, BMI: body mass index, BS: bariatric surgery*

- A. Respondents with \geq WHO Class II Obesity vs. Respondents with $<$ WHO Class II Obesity (= reference). Based on the entire cohort [$n = 1482$; $R^2 = 0.156$]
- 340 B. Respondents who underwent bariatric surgery vs. Respondents without history of bariatric surgery (= reference). Based on the entire cohort [$n = 1482$, $R^2 = 0.169$]
- C. Respondents with an interest to undergo bariatric surgery vs. Respondents not interested to undergo bariatric surgery (= reference). Within the subgroup of respondents without history of bariatric surgery [$n = 409$; $R^2 = 0.142$]

345

Figures

350 **Figure 1.** Free-word association networks to “weight loss” in patients with obesity. The modules reflect the data-driven frequency of co-occurrences of associations. Potential interpretation: *Green: benefit-focused mindset; Purple: procedure-focused mindset.*

Figure 2. Frequency of emotional labels assigned to the associations of each cognitive module

355

Supplemental Material

Supplemental Methods

Supplemental Method 1. The survey as it was presented to the participants (MethodS1.pdf)

Supplemental Table

360 **Supplemental Table 1.**

A. Descriptive characteristics of the respondents based on their current stage of obesity and on their bariatric surgery status. *BMI: body mass index*

B. Comparison of demographic parameters between subgroups

365 **Supplemental Figures**

Supplemental Figure 1. Flowchart of the selection of the study cohort

Supplemental Figure 2. Attractive and repulsive associations

Supplemental Figure 3. Bridging associations

370 **Supplemental Figure 4.** Correlation of current BMI and Body Image Dissatisfaction

References

- [1] Bray GA, Fruhbeck G, Ryan DH, Wilding JP. Management of obesity. *Lancet*. 2016;387:1947-56.
- [2] Kaplan LM, Golden A, Jinnett K, Kolotkin RL, Kyle TK, Look M, et al. Perceptions of Barriers to Effective Obesity Care: Results from the National ACTION Study. *Obesity (Silver Spring)*. 2018;26:61-9.
- [3] Puhl RM, Heuer CA. Obesity stigma: important considerations for public health. *Am J Public Health*. 2010;100:1019-28.
- [4] Gero D, Hulesch B, Bueter M. Mindset and Communication Barriers in the Diffusion of Bariatric Surgery. *Curr Atheroscler Rep*. 2018;20:38.
- [5] Munoz DJ, Lal M, Chen EY, Mansour M, Fischer S, Roehrig M, et al. Why patients seek bariatric surgery: a qualitative and quantitative analysis of patient motivation. *Obes Surg*. 2007;17:1487-91.
- [6] Libeton M, Dixon JB, Laurie C, O'Brien PE. Patient motivation for bariatric surgery: characteristics and impact on outcomes. *Obes Surg*. 2004;14:392-8.
- [7] Edward KL, Hii MW, Giandinoto JA, Hennessy J, Thompson L. Personal Descriptions of Life Before and After Bariatric Surgery From Overweight or Obese Men. *Am J Mens Health*. 2018;12:265-73.
- [8] Peacock JC, Perry L, Morien K. Bariatric patients' reported motivations for surgery and their relationship to weight status and health. *Surg Obes Relat Dis*. 2018;14:39-45.
- [9] Sharman MJ, Venn AJ, Hensher M, Wilkinson S, Palmer AJ, Williams D, et al. Motivations for Seeking Bariatric Surgery: The Importance of Health Professionals and Social Networks. *Bariatr Surg Pract P*. 2016;11:104-9.
- [10] Walach H, Loughlin M. Patients and agents - or why we need a different narrative: a philosophical analysis. *Philos Ethics Humanit Med*. 2018;13:13.
- [11] Teutsch C. Patient-doctor communication. *Med Clin North Am*. 2003;87:1115-45.
- [12] File B, Keczer Z, Vancso A, Bothe B, Toth-Kiraly I, Hunyadi M, et al. Emergence of polarized opinions from free association networks. *Behav Res Methods*. 2018.
- [13] Luduena GA, Behzad MD, Gros C. Exploration in free word association networks: models and experiment. *Cogn Process*. 2014;15:195-200.
- [14] Vijayasarithi A, Kharkar R, Salamon N. Strategies for Patient-Centered Communication in the Digital Age. *Curr Probl Diagn Radiol*. 2018.
- [15] Eysenbach G. Improving the quality of Web surveys: the Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res*. 2004;6:e34.
- [16] Ekman P. Are there basic emotions? *Psychol Rev*. 1992;99:550-3.
- [17] Robinson DL. Brain function, emotional experience and personality. *Netherlands Journal of Psychology*. 2008;64:152-68.
- [18] Blanch L. Under Construction: Body Image and Self-Representation in Virtual Worlds. . Toronto, Canada: York University, Toronto, Canada; 2014.
- [19] Ratcliff MB, Eshleman KE, Reiter-Purtill J, Zeller MH. Prospective changes in body image dissatisfaction among adolescent bariatric patients: the importance of body size estimation. *Surg Obes Relat Dis*. 2012;8:470-5.
- [20] Research AAFPO. Standard Definitions: Final Dispositions of Case Codes and Outcome Rates for Surveys. 8th ed. Oakbrook Terrace, IL, USA2015.
- [21] Lancichinetti A, Fortunato S. Consensus clustering in complex networks. *Sci Rep*. 2012;2:336.
- [22] Rozier MD, Ghaferi AA, Rose A, Simon NJ, Birkmeyer N, Prosser LA. Patient Preferences for Bariatric Surgery: Findings From a Survey Using Discrete Choice Experiment Methodology. *JAMA Surg*. 2018:e184375.
- [23] Mondragon NI, Txertudi MB. Understanding fatness in the public sphere of young students: social representations and emotional response. *Cad Saude Publica*. 2018;34:e00197917.
- [24] Torelli CJ, Kaikati AM. Values as predictors of judgments and behaviors: the role of abstract and concrete mindsets. *J Pers Soc Psychol*. 2009;96:231-47.

420 [25] Oettingen G, Wadden TAJCT, Research. Expectation, fantasy, and weight loss: Is the impact of positive thinking always positive? 1991;15:167-75.

[26] Trottier K, Polivy J, Herman CP. Effects of resolving to change one's own behavior: expectations vs. experience. *Behav Ther.* 2009;40:164-70.

[27] Monpellier VM, Antoniou EE, Mulkens S, Janssen IMC, van der Molen ABM, Jansen ATM. Body image dissatisfaction and depression in postbariatric patients is associated with less weight loss and a desire for

425 body contouring surgery. *Surg Obes Relat Dis.* 2018;14:1507-15.

[28] Gallagher MWCJ. *The Oxford handbook of hope.* New York, NY: Oxford University Press; 2018.

[29] McCullough ME, Kilpatrick SD, Emmons RA, Larson DB. Is gratitude a moral affect? *Psychol Bull.* 2001;127:249-66.

[30] Bollo H, Bothe B, Toth-Kiraly I, Orosz G. Pride and Social Status. *Front Psychol.* 2018;9:1979.

430 [31] Carvalho A, Jr., Turato ER, Chaim EA, Magdaleno R, Jr. Weight regain among women after metabolic and bariatric surgery: a qualitative study in Brazil. *Trends Psychiatry Psychother.* 2014;36:140-6.

[32] Karagulle OO, Yavuz E, Gulcicek OB, Solmaz A, Senturk S, Erdogan A, et al. Psychological well-being and marital satisfaction in response to weight loss after bariatric surgery. *Surg Today.* 2019.

[33] Conceição EM, de Lourdes M, Moreira L, Pinto-Bastos A, Félix S. Weight loss expectations and weight loss

435 after surgery: the mediating role of body image and weight concerns. *Surgery for Obesity and Related Diseases.* 2020.

[34] Sharma AM, Belanger A, Carson V, Krah J, Langlois MF, Lawlor D, et al. Perceptions of barriers to effective obesity management in Canada: Results from the ACTION study. *Clin Obes.* 2019;9:e12329.

[35] Aij KH, Teunissen M. Lean leadership attributes: a systematic review of the literature. *J Health Organ*

440 *Manag.* 2017;31:713-29.

[36] Teichert TA, Schöntag K. Exploring consumer knowledge structures using associative network analysis. 2010;27:369-98.

[37] Powell SM, Fasczewski KS, Gill DL, Davis PG. Go with the FLOW: Implementation of a psychological skills intervention in an exercise program for post-bariatric surgery patients. *J Health Psychol.*

445 2018;1359105318793182.

[38] Nowicki T, Burns C, Fulbrook P, Jones J. Changing the mindset: an inter-disciplinary approach to management of the bariatric patient. *Collegian.* 2009;16:171-5.

[39] Vallone D, Greenberg M, Xiao H, Bennett M, Cantrell J, Rath J, et al. The Effect of Branding to Promote Healthy Behavior: Reducing Tobacco Use among Youth and Young Adults. *Int J Environ Res Public Health.*

450 2017;14.

[40] Neermark S, Holst C, Bisgaard T, Bay-Nielsen M, Becker U, Tolstrup JS. Validation and calibration of self-reported height and weight in the Danish Health Examination Survey. *Eur J Public Health.* 2018.

Tables

Table 1. Respondents' characteristics according to cognitive module membership. *BMI: Body Mass Index. BS: bariatric surgery. *: significant ($P < 0.01$) difference in the frequency of the emotion between the two modules.*

	Module of associations		P
	Benefit-focused	Procedure-focused	
n	1174	253	
Age (mean, SD)	43.1 ± 10.8	40.4 ± 10.2	< 0.001
Female (%)	94.5	92.5	1
Relationship status (%)			0.22
<i>single</i>	15	17	
<i>divorced</i>	11	8	
<i>in relation</i>	22	26	
<i>married</i>	50	48	
<i>widow</i>	2	2	
Current BMI ≥35 kg/m² (%)	54.1	43.1	0.002
Maximal lifetime BMI (mean, SD)	50.7 ± 8.5	50.3 ± 9.2	0.51
Body Image Dissatisfaction (mean, SD)	2.3 ± 1.7	2.2 ± 1.5	0.52
Proportion of positive emotions (%)	86 ± 19	47 ± 27	< 0.001
Had bariatric surgery (%)	71.7	76.3	0.14
Among those who did not have BS:			
<i>Interested in undergoing BS (%)</i>	87	66.7	< 0.001
Most frequent associations (%)			
1 st	health (54)	sport (17)	
2 nd	agility (24)	diet (14)	
3 rd	attractiveness (24)	effort (9)	
4 th	happiness (21)	excess skin (5)	
5 th	activity (21)	sacrifice (4)	
Most frequently associated emotions (%)			
1 st	joy (21) *	hope (13) *	
2 nd	gratitude (17) *	frustration (13) *	
3 rd	hope (17) *	anxiety (11) *	
4 th	pride (12) *	interest (9)	
5 th	interest (7)	joy (7) *	

Table 2. Explorative models using logistic regression investigating the effect of the stage of obesity and of bariatric surgery status on cognitive module membership and use of different emotional labels. *CI: confidence interval, BMI: body mass index BS: bariatric surgery*

A. Respondents with \geq WHO Class II Obesity vs. Respondents with $<$ WHO Class II Obesity (= reference). Based on the entire cohort [$n = 1482$; $R^2 = 0.156$]

	Respondents with \geq Class II obesity			
	<i>Odds ratio</i>	<i>95% CI</i>		<i>P</i>
Benefit-focused cognitive module	2.2	1.6	3.1	<0.001
Joy	1.4	0.5	4.2	0.5
Hope	76.6	25.7	227.8	<0.001
Gratitude	0.04	0.01	0.1	<0.001
Pride	0.2	0.06	0.7	0.01
Interest	0.4	0.1	1.4	0.14
Anxiety	0.7	0.2	4.9	0.96
Surprise	0.2	0.04	0.9	0.04
Frustration	1.6	0.3	9.4	0.63

B. Respondents who underwent bariatric surgery vs. Respondents without history of bariatric surgery (= reference). Based on the entire cohort [$n = 1482$, $R^2 = 0.169$]

	Respondents with prior bariatric surgery			
	<i>Odds ratio</i>	<i>95% CI</i>		<i>P</i>
Benefit-focused cognitive module	0.4	0.3	0.6	<0.001
Joy	0.5	0.2	1.8	0.3
Hope	0.03	0.01	0.1	<0.001
Gratitude	107.8	27.7	420.2	<0.001
Pride	15.6	3.4	69.2	<0.001
Interest	1	0.3	3.5	0.99
Anxiety	0.6	0.1	3.6	0.6
Surprise	7.2	1.2	45	0.04
Frustration	0.2	0.03	1.2	0.08

- C. Respondents with an interest to undergo bariatric surgery vs. Respondents not interested to undergo bariatric surgery (= reference). Within the subgroup of respondents without history of bariatric surgery [n = 409; R² = 0.142]

	Respondent with an interest to undergo bariatric surgery			
	<i>Odds ratio</i>	<i>95% CI</i>		<i>P</i>
Benefit-focused cognitive module	<i>2.4</i>	<i>1.17</i>	<i>4.94</i>	<i>0.02</i>
Joy	0.9	0.06	15.2	0.96
Hope	<i>142.5</i>	<i>9.9</i>	<i>2047.1</i>	<i><0.001</i>
Gratitude	1.2	0.06	26	0.9
Pride	0.3	0.01	9.2	0.5
Interest	0.1	0.01	1.8	0.12
Anxiety	5.3	0.1	276.7	0.41
Surprise	4.3	0.06	328.7	0.51
Frustration	0.8	0.02	47	0.93






Figure 2.



Gero, File, et al. Mental and emotional representations of “weight loss”: free-word association networks in members of bariatric surgery-related social media communities. SURGERY FOR OBESITY AND RELATED DISEASES. 2019

Highlights


- Aim to map mental and emotional representation of *weight loss* in bariatric patients
- 1st application of a novel network-analysis based methodology in the context of BS
- 1482 patients were surveyed within bariatric social media communities
- Patients think about weight loss in two cognitive schemes: benefit/process focused
- Benefit-focused respondents were more likely to consider bariatric surgery
- Novel communication strategies may incorporate the patients’ free associations



[Click here to access/download](#)

Supplementary Materials

WeightlossNetworks_SOARD_Supplement_R2V2.docx





Click here to access/download
Supplementary Materials
MethodsS1.pdf